

Constraints on $\tan\beta$ in SUSY SU(5) GUT

Jun Tabei*

Department of Physics, Waseda University, Tokyo 169, Japan

Hiroshi Hotta†

*Institute of Material Science and Technology,
Waseda University, Tokyo 169, Japan*

(Dated: February 1, 2008)

Abstract

On the evaluation of $\tan\beta$ is revisited by the method of the renormalization group with the criteria of the bottom-tau unification in the SUSY SU(5) GUT model. Among the conventional supersymmetric theories, an energy-scale constant M_{SUSY} is introduced by-hand to decouple the lower energy-scale region from the effects of the supersymmetric particles. Provided that M_{SUSY} really exists, only small $\tan\beta \simeq 2$ is allowed theoretically. While, medium value as $\tan\beta = 6 \sim 25$ is also supported if M_{SUSY} is vanished away from the model.

PACS numbers: 12.60.Jv, 12.10.-g, 11.10.Hi

*Electronic address: jun@hep.phys.waseda.ac.jp

†Electronic address: hotta@hep.phys.waseda.ac.jp

I. INTRODUCTION

1) - What are unified ?

As one of the general features of the SU(5) GUT models, both Yukawa coupling constants of the bottom-quark and the tau-lepton are unified at GUT scale. Additionally, in several SU(5) models including the *SUSY* SU(5) GUT model[1], the evidence for the unification of gauge coupling constants is numerically provided[2] .

2) - What is M_{SUSY} ?

In the conventional analysis[2], the gauge unification in the *SUSY* SU(5) GUT model is achieved by introducing an energy-scale constant M_{SUSY} by-hand to decouple the lower energy-scale region from the effects of the supersymmetric particles. Thus, M_{SUSY} is regarded as the threshold between the standard model(SM) and *SUSY*. The value of M_{SUSY} depends on the strong coupling constant α_3 very sensitively[2]. For instance, while $M_{SUSY} \simeq 1(\text{TeV})$ for $\alpha_3 = 0.106$ is given[2] in 1991, $M_{SUSY} \leq 100(\text{GeV})$ is obtained for the recent value[3] of $\alpha_3 = 0.1185$. If M_{SUSY} were really less than 100(GeV), we should have already detected such supersymmetric particles in the region of $\sim 100(\text{GeV})$ experimentally, however we have not yet. This fact may allow us to think of M_{SUSY} as a negligible parameter, or, at least, the authors should discuss on the possibility of the models without M_{SUSY} . Note that such model without M_{SUSY} means the pure supersymmetric theory without including the conventional standard model.

3) How much $\tan \beta$ is optimal ?

$\tan \beta$ is evaluated by the two different ways of the likelihood analysis at two-loop level in this paper. One of the two ways is on the unification of the gauge coupling constants, and another is of the bottom-tau's Yukawa coupling constants. The sequence to optimize $\tan \beta$ is as follows:

In the first, M_{SUSY} and the GUT scale M_X are fixed by the χ^2 -fitting on the unification of the gauge(χ_g^2) coupling constants. Once M_{SUSY} and M_X are fixed, corresponding $\tan \beta$ is derived numerically to realize the bottom-tau unification at M_X scale.

Additionally, the analysis on the model without M_{SUSY} is also carried out in the same manner.

II. ANALYSIS OF $\tan\beta$ WITH M_{SUSY}

On the optimization of $\tan\beta$ is discussed[4] by the renormalization group equations(RGEs) of the gauge and Yukawa coupling constants at two-loop level[5, 6] in the SM and SUSY. As the Yukawa coupling constants, heavier three of the ordinal Fermions (top-quark, bottom-quark, and tau-lepton) are taken into consideration.

First of all, the explicit values of the parameters in this paper are referred from the latest issue of the Particle Data Group[3].

On the boundary conditions of the RGEs, they are described further below. The conditions of the gauge coupling constants are given on Z-boson's mass shell $M_Z = 91.184(\text{GeV})$ in the $\overline{\text{MS}}$ -scheme as:

$$\alpha_1 = \frac{5}{3} \frac{\alpha_{\overline{\text{MS}}}}{1 - \sin^2 \theta_{\overline{\text{MS}}}} , \quad (1)$$

$$\alpha_2 = \frac{\alpha_{\overline{\text{MS}}}}{\sin^2 \theta_{\overline{\text{MS}}}} , \quad (2)$$

as their explicit values are:

$$\alpha_{\overline{\text{MS}}}^{-1} = 127.934 \pm 0.027 , \quad (3)$$

$$\sin^2 \theta_{\overline{\text{MS}}} = 0.23117 \pm 0.00016 , \quad (4)$$

$$\alpha_3 = 0.1185 \pm 0.0020 . \quad (5)$$

Moreover, the definition of $\tan\beta$ at M_Z is given as usual:

$$\tan\beta(M_Z) \doteq v_2(M_Z)/v_1(M_Z) , \quad (6)$$

where, v_1 and v_2 imply the vacuum expectation values(VEVs) of two Higgs doublets, and as their explicit value:

$$v(M_Z) = (v_1^2(M_Z) + v_2^2(M_Z))^{1/2} = 246(\text{GeV}) , \quad (7)$$

is fixed. For simplicity, mere $\tan\beta$ means $\tan\beta(M_Z)$ in this paper.

On the Higgs quartic coupling constant λ , this is precisely defined at M_{SUSY} [4] as:

$$\lambda(M_{SUSY}) = \pi \left[\frac{3}{5} \alpha_1(M_{SUSY}) + \alpha_2(M_{SUSY}) \right] \cos^2 2\beta(M_{SUSY}) , \quad (8)$$

however, it is very difficult to derive λ from this definition. As λ , the authors make use of an approximate, and easier alternative definition at M_Z :

$$\lambda \doteq \pi \left[\frac{3}{5} \alpha_1(M_Z) + \alpha_2(M_Z) \right] \cos^2 2\beta(M_Z) . \quad (9)$$

Note that the numerical difference between the definitions Eq. (8) and Eq. (9) is very small as about 0.1%, and λ is decoupled to the Yukawa coupling constants among the RGEs at one-loop level, therefore, all the numerical results in this paper are effectually independent of the choice of the definition of λ .

Boundary conditions of the Yukawa coupling constants are given at each on-shell energy-scales as:

$$\alpha_q(m_q) = \frac{m_q^2(m_q)}{2\pi v^2(m_q)} , \quad (10)$$

where, the suffix q implies the flavor of Fermions as $q = t, b$, or τ , respectively. Their explicit values are as follows:

$$m_t(m_t) = 174.3 \pm 5.1(\text{GeV}) , \quad (11)$$

$$m_b(m_b) = 4.2 \pm 0.2(\text{GeV}) , \quad (12)$$

$$m_\tau(m_\tau) = 1.77699 \pm 0.00029(\text{GeV}) . \quad (13)$$

When assuming the existence of M_{SUSY} , corresponding boundary conditions are given as:

$$\alpha_i^{SUSY}(M_{SUSY}) = \alpha_i^{SM}(M_{SUSY}) , \quad (i = 1, 2, 3) , \quad (14)$$

$$\begin{aligned} \alpha_t^{SUSY}(M_{SUSY}) &= \frac{m_t^2(M_{SUSY})}{2\pi v^2(M_{SUSY}) \sin^2 \beta(M_{SUSY})} \\ &= \alpha_t^{SM}(M_{SUSY}) \frac{1}{\sin^2 \beta(M_{SUSY})} \\ &= \alpha_t'^{SM}(M_{SUSY}) , \end{aligned} \quad (15)$$

$$\begin{aligned} \alpha_{b,\tau}^{SUSY}(M_{SUSY}) &= \alpha_{b,\tau}^{SM}(M_{SUSY}) \frac{1}{\cos^2 \beta(M_{SUSY})} \\ &= \alpha_{b,\tau}'^{SM}(M_{SUSY}) , \end{aligned} \quad (16)$$

where, the variables with a prime(') mean the redefinition of them in the SM region to keep their continuity between the regions of the SM and SUSY. With such redefinitions, the graphs in this paper is continuous even if crossing on the M_{SUSY} border.

The conditions at GUT energy-scale M_X are described in the following subsections.

A. On the gauge coupling constants

The average value α_g among α_1, α_2 , and α_3 is introduced around the GUT scale M_X . χ_g^2 -fitting function is defined as the squared sum of each differences between α_g and $\alpha_i (i = 1 \sim 3)$ as follows:

$$\chi_g^2 \doteq \sum_{i=1}^3 \left(\frac{\alpha_i - \alpha_g}{\delta\alpha_i} \right)^2, \quad (17)$$

where, $\delta\alpha_i$ means the uncertainty width of α_i , respectively. the GUT scale M_X is defined as the energy-scale where χ_g^2 becomes the minimum. In order to make χ_g^2 the minimum, M_{SUSY} is fine-tuned. As the results, we find $\chi_g^2 \simeq 0$ can be achieved by tuning M_{SUSY} at any $\tan\beta$ in the ranges of the errorbars of the other parameters. Fine-tuned values of M_{SUSY} are shown as functions of $\tan\beta$ with the uncertainty width of α_3 in Fig. 1. We find the derived M_{SUSY} 's value depends on only the strong coupling constant α_3 rather than the other parameters.

B. On the Yukawa coupling constants

When the SU(5) GUT scenario is presumed, Yukawa coupling constants of the bottom-quark and tau-lepton are unified from each other at M_X . Their χ_Y^2 -fitting function is defined as the following equations:

$$\chi_Y^2 \doteq \sum_{q=b,\tau} \left(\frac{\alpha_q - \alpha_Y}{\delta\alpha_q} \right)^2, \quad (18)$$

$$\alpha_Y = (\alpha_b + \alpha_\tau) / 2, \quad (19)$$

where, q implies b or τ , and $\delta\alpha_q$ means the uncertainty width of α_q , respectively. With the uncertainty width $\delta = 0.2(\text{GeV})$ of m_b , χ_Y^2 at M_X scale as functions of $\tan\beta$ are shown in Fig. 2. The behavior of χ_Y^2 is not sensitive on the variance of m_t or α_3 . Also shown in Fig. 2, only small $\tan\beta \simeq 2$ is allowed by the likelihood analysis with the χ_Y^2 function, and this minimal point of χ_Y^2 is stable despite of the uncertainty of the parameters like m_b , m_t , or α_3 .

III. ANALYSIS OF $\tan\beta$ WITHOUT M_{SUSY}

As mentioned above, on the model without M_{SUSY} is discussed in this section. Therefore, the RGEs[6] in this section are pure SUSY. On the boundary conditions of the gauge coupling constants, the VEVs of Higgs bosons, and $\tan\beta$, they are given at M_Z in the same way of the model with including M_{SUSY} as described in the previous section as from Eq. (3) to Eq. (6). The boundary conditions of the Yukawa coupling constants are given at the energy-scales of each on-shell masses as follows:

$$\alpha_t(m_t) = \frac{m_t^2(m_t)}{2\pi v^2(m_t) \sin^2 \beta(m_t)} , \quad (20)$$

and

$$\alpha_{b,\tau}(m_{b,\tau}) = \frac{m_{b,\tau}^2(m_{b,\tau})}{2\pi v^2(m_{b,\tau}) \cos^2 \beta(m_{b,\tau})} . \quad (21)$$

The definitions of χ_g^2 and χ_Y^2 at M_X are the same ones Eq. (17) and Eq. (18) in the previous section, however, in this section, χ_g^2 depends on $\tan\beta$ or other parameters directly in contrast to the previous section, because the decoupling boundary M_{SUSY} is vanished away from the model.

The dependence of $\tan\beta$ on χ_g^2 with the uncertainty width of α_3 is shown in Fig. 3 . As Fig. 3 shows, two minimal points exist at small and large $\tan\beta$. The variance of χ_Y^2 at M_X is shown with each uncertainty width of m_b , m_τ , or α_3 in Fig. 4a, 4b, or 4c, respectively. As shown in these graphs, usually χ_Y^2 has three local minima.

IV. DISCUSSIONS

A. On the model with M_{SUSY}

Since χ^2 of the gauge coupling constants(χ_g^2) can be always equal to zero by fine-tuning of M_{SUSY} , there exists no effective constraint on $\tan\beta$ from the unification of the gauge coupling constants. $\tan\beta$ is evaluated as 1.6 by the criteria of the bottom-tau unification in this model with M_{SUSY} . M_{SUSY} is estimated about 2(TeV) to unify the gauge coupling constants at two-loop level. (As a reference, $M_{SUSY} \ll M_Z$ is derived at one-loop level.) Nevertheless, such small $\tan\beta \simeq 2$ is difficult to be allowed by $g-2$ experiments[7], because small $\tan\beta$ makes the masses of sparticles quite smaller than expected. The evolutions

of the gauge and Yukawa coupling constants at $\tan\beta = 1.6$ are shown in Fig. 6a and 6b, respectively. The summary of the arguments in this subsection is shown in the Table I.

B. On the model without M_{SUSY}

The value of $\tan\beta$ is possibly settled at 2, $6 \sim 25$, or 50 in accordance with three minima of χ_Y^2 . However, large $\tan\beta \simeq 50$ is denied theoretically, because χ_g^2 and χ_Y^2 cannot be minimal simultaneously, *i.e.*, they are exclusive from each other as shown in Fig. 5. On the small $\tan\beta \simeq 2$ choice, both of χ_g^2 and χ_Y^2 are enough small, however, this is excluded by the recent $g - 2$ experiments[7] as mentioned previously. Large $\tan\beta \simeq 50$ is also not probable if $b \rightarrow s\gamma$ experiment[8] is true. Moreover, the authors are dubious whether these two local minima of χ_Y^2 at small and large $\tan\beta$ are physical ones or not, because they make the RGEs divergent at one-loop level, therefore, the evolution of the Yukawa coupling constants seems strange even at two-loop level, for instance, as shown in Fig. 7 at $\tan\beta = 1.9$. Medium $\tan\beta$ as $6 \sim 25$ is good for χ_Y^2 of the bottom-tau unification, however, χ_g^2 of the gauge unification cannot be enough small at two-loop level. Thus, precise unification of the gauge coupling constants cannot occur with medium $\tan\beta$ as $6 \sim 25$. The evolutions of the gauge or Yukawa coupling constants at $\tan\beta = 6.0$ are shown in Fig. 8a or 8b, respectively. These discussions are summarized in the Table II.

V. SUMMARY AND CONCLUSION

In the SU(5) GUT scenario, $\tan\beta$ is evaluated by making use of the RGEs of the gauge and Yukawa coupling constants. Only small $\tan\beta \simeq 2$ is allowed in the model including M_{SUSY} , and $\tan\beta$ is estimated as 2 or 6 to 25 in the model without M_{SUSY} , theoretically. However, small $\tan\beta \simeq 2$ is difficult to be allowed by the analysis of the recent $g - 2$ experiment[7]. As the result, only medium $\tan\beta$ as $6 \sim 25$ is acceptable in the SUSY SU(5) GUT model without M_{SUSY} if the bottom-tau unification is prior to the unification among the gauge coupling constants.

Acknowledgments

The authors thank Y. Okada, M. Kakizaki and T. Nihei for useful advice. We are grateful to D. W. Hertzog for helpful comments on our preliminary stage.

- [1] H. P. Nilles, Phys. Rep. **110**, 1(1984)
- [2] U. Amaldi, W. de Boer, and H. Fürstenau, Phys. Lett. **B260**, 447(1991)
- [3] The Particle Data Group, browse on <http://pdg.lbl.gov/>
- [4] V. Barger, M. S. Berger, and P. Ohmann, Phys. Rev. **D47**, 1093(1993)
- [5] H. Arason, *et al.*, Phys. Rev. **D46**, 3945(1992)
- [6] D. J. Castaño, E. J. Piard, and P. Ramond, Phys. Rev. **D49**, 4882(1994)
- [7] As the original report of the precise $g - 2$ measurement:
H. N. Brown, *et al.*(Muon ($g - 2$) Collaboration), Phys. Rev. Lett. **86**, 2227(2001) ;
as the theoretical analysis:
U. Chattopadhyay and P. Nath, Phys. Rev. Lett. **86**, 5854(2001)
- [8] G. C. Cho, *et al.*, hep-ph/9901351 ;
H. Baer, *et al.*, Phys. Rev. **D58**, 015007(1998)

TABLE I: Possiblity of $\tan \beta$ with M_{SUSY} .

$\tan \beta$	small($\simeq 2$)	medium($6 \sim 25$)	large($\simeq 50$)
χ_g^2	○	○	○
χ_Y^2	○	×	×
$g - 2$	×	○	○

TABLE II: Possiblity of $\tan \beta$ without M_{SUSY} .

$\tan \beta$	small($\simeq 2$)	medium($6 \sim 25$)	large($\simeq 50$)
χ_g^2	○	Δ	$\circ(\times)^a$
χ_Y^2	○	○	$\times(\circ)^a$
$g - 2$	×	○	○
$b \rightarrow s\gamma$	○	○	Δ

^aThese two scores are exclusive from each other.




Fig. 1

FIG. 1: M_{SUSY} as functions of $\tan\beta$ for $\alpha_3 \pm \delta = 0.1185 \pm 0.0020$.




Fig. 2

FIG. 2: χ^2 of the Yukawa coupling constants(χ_Y^2) at GUT scale M_X for $m_b \pm \delta = 4.2 \pm 0.2(\text{GeV})$ with M_{SUSY} .




Fig. 3

FIG. 3: χ^2 of the gauge coupling constants(χ^2_g) at M_X for $\alpha_3 \pm \delta = 0.1185 \pm 0.0020$ without M_{SUSY} .




Fig. 4a




Fig. 4b




Fig. 4c

FIG. 4: χ_Y^2 at M_X for each uncertainty width of (a) m_b , (b) m_t , and (c) α_3 without M_{SUSY} , respectively.




Fig. 5

FIG. 5: Opposite behavior of χ_Y^2 vs χ_g^2 around large $\tan\beta \simeq 50$ at M_X without M_{SUSY} .




Fig. 6a




Fig. 6b

FIG. 6: The evolutions of the (a)gauge and (b)Yukawa coupling constants at $\tan\beta=1.6$ with M_{SUSY} .




Fig. 7

FIG. 7: The evolution of the Yukawa coupling constants at $\tan\beta=1.9$ without M_{SUSY} .




Fig. 8a




Fig. 8b

FIG. 8: The evolutions of the (a)gauge and (b)Yukawa coupling constants at $\tan\beta=6.0$ without M_{SUSY} .